

PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM B

Search PubMed for [ ] Go Clear  
Limits Preview/Index History Clipboard Details

About Entrez

Text Version

Display Abstract Show: 20 Sort Send to Text

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: Nature 1978 Dec 21-28;276(5690):785-90

Related Articles, Li

## Rearrangement of genetic information may produce immunoglobulin diversity.

Weigert M, Gatmaitan L, Loh E, Schilling J, Hood L.

The nearly complete amino-acid sequences of 22 closely related immunoglobulin kappa variable (Vkappa) regions from the inbred NZB mouse are presented. This group of Vkappa regions is encoded by at least six germline Vkappa genes. These data also suggest that the mouse kappa gene is divided into three segments termed V or variable (residues 1 to 98 or 99), J or joining (residues 99 or 100 to 112) and C or constant (residues 113--219). Tonegawa et al. have recently described a similar J segment for mouse lambda chains. Inbred mice contain multiple Vkappa and Jkappa gene segments. Therefore, different combinations of V and J gene segments may be joined at the DN level during the differentiation of individual lymphocytes to contribute to antibody diversity.

PMID: 103003 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text


Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer

Mar 17 2003 10:14



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

B

Search

PubMed

for

CD30 antibody AND chemotherapy

Preview

Go

C

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

- Search History will be lost after one hour of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.

Search

Most Recent Queries

Time Result

#5 Search CD30 antibody AND chemotherapy

08:36:28

55

#9 Related Articles for PubMed (Select 10845927)

08:34:14

287

#7 Related Articles for PubMed (Select 12402667)

08:29:52

101

#3 Related Articles for PubMed (Select 12560434)

08:26:08

118

#1 Search her2 AND chemotherapy

08:25:32

861

Clear History

Write to the Help Desk

NCBI | NLM | NIH

Department of Health &amp; Human Services

Freedom of Information Act | Disclaimer

Mar 17 2003 10:44:01

[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[PMC](#)[Taxonomy](#)[OMIM](#)[B](#)

Search

PubMed

for

Go

Clear

[Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)[About Entrez](#)

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

[Overview](#)[Help | FAQ](#)[Tutorial](#)[New/Noteworthy](#)[E-Utilities](#)

PubMed Services

[Journals Database](#)[MeSH Browser](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[LinkOut](#)[Cubby](#)

Related Resources

[Order Documents](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)[Privacy Policy](#)☐ 1: Cancer Res 1995 Nov 15;55(22):5335-41[Related Articles, Li](#)


### **In vivo antitumor effects of unconjugated CD30 monoclonal antibodies on human anaplastic large-cell lymphoma xenografts.**

Tian ZG, Longo DL, Funakoshi S, Asai O, Ferris DK, Widmer M, Murphy WJ.

Laboratory of Leukocyte Biology, National Cancer Institute-Frederick Cancer Research and Development Center, Maryland 21702, USA.

CD30 is a M(r) 120,000 surface antigen identified originally by the Ki-1 monoclonal antibody (moAb) against primary and cultured Reed-Sternberg cells present in Hodgkin's disease and anaplastic large-cell lymphomas (ALCLs). Examination of two ALCL cell lines (Karpas 299 and Michel) demonstrated cell surface expression of CD30. Incubation of these lymphomas with two anti-CD30 moAbs that recognize the ligand-binding site (M44 or HeFi-1) resulted in significant growth inhibition in vitro with significant decreases in cell viability. Another anti-CD30 moAb, Ber-H2, which recognizes a determinant not involved in ligand binding, had no effect on ALCL growth in vitro. When these human ALCL lines were transferred i.v. into mice with severe combined immune deficiency, the mice developed extensive metastasis in the s.c., brain, or eye tissues. The treatment of mice with either M44 or HeFi-1 anti-CD30 moAbs resulted in significant increases in survival, with some mice remaining disease free for more than 100 days. Thus, anti-CD30 treatment is efficacious for CD30+ ALCL cell lines in vivo, and unconjugated anti-CD30 moAbs may be of potential clinical use.

PMID: 7585597 [PubMed - indexed for MEDLINE]



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

B

Search

PubMed

for

Limits

Preview/Index

History

Clipboard

Details

Go

Clear

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Nature 1978 Dec 21-28;276(5690):785-90

Related Articles, Li

## Rearrangement of genetic information may produce immunoglobulin diversity.

Weigert M, Gatmaitan L, Loh E, Schilling J, Hood L.

The nearly complete amino-acid sequences of 22 closely related immunoglobulin kappa variable (Vkappa) regions from the inbred NZB mouse are presented. This group of Vkappa regions is encoded by at least six germline Vkappa genes. These data also suggest that the mouse kappa gene is divided into three segments termed V or variable (residues 1 to 98 or 99), J or joining (residues 99 or 100 to 112) and C or constant (residues 113--219). Tonegawa et al. have recently described a similar J segment for mouse lambda chains. Inbred mice contain multiple Vkappa and Jkappa gene segments. Therefore, different combinations of V and J gene segments may be joined at the DN level during the differentiation of individual lymphocytes to contribute to antibody diversity.

PMID: 103003 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to

Text

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

Mar 17 2003 10:44